

Use of a Miami J Collar following an unstable neck injury or fracture



Important!

2 people are required to remove or apply the collar.

This leaflet is designed to provide patients with information on a Miami J collar, and to help their carers, family or friends understand how to safely remove and apply the collar as they cannot do this themselves.



Use of the Miami J Collar:

Following a cervical (neck) spinal injury it is important to prevent excess movement of the spine in order to allow the fracture or damaged area to heal.

We can help your neck to heal and reduce the chance of further injury or nerve damage by applying a neck collar. This neck collar is called a Miami J Collar.

Aim of the Miami J Collar:

You wear the collar as an external splint to protect your neck against excessive flexion (bending forward) and rotation (twisting) while the fracture or damage is healing.

When does someone start wearing the collar?

After injury, if a collar is required, a Healthcare Professional will measure and supply you with the correct size of collar and check it fits properly.

When should the collar be worn?

The collar should be worn at all times other than when your neck is being cleaned or the pads are being changed. This needs to be carried out by 2 people.

Pictures of Miami J Collar



Front of Collar

Back of Collar



Collar with pads removed

Important!

2 people are required to apply or remove the collar.

Applying and removing the Miami J Collar

A video to assist with the fitting of the collar can be found here:

Miami J Collar Unstable 2 Person Fitting Video

▶ <https://youtu.be/LQILsi70f-c>




Or alternatively, written instructions are below.

The person wearing the Miami J Collar should lie flat on their back. One person should firmly hold the head in a neutral position. This same person should hold the head until the collar is securely fitted again.

To put the collar on, the second person should take the back section of the collar (the word "back" is written in the plastic and an arrow indicates the correct way up). Position the back section behind your neck by pushing it down into the bed or thin pillow and under the neck following its curve until it's central (See Picture 1).

Picture 1



When the back section is in place, take the front section of the collar and scoop this up underneath the chin and jaw, making sure it contacts evenly, pushing the collar upwards and backwards in the direction of the ears (See Picture 2). The white plastic sections of the front and back of the collar do not need to overlap, as long as they meet. The chin should be held in the centre of the chin piece. Make sure the collar is central by checking there are equal amounts of plastic on each side.

Picture 2



Fasten one strap at a time while holding the collar securely at the front, making sure the collar remains centred. The straps come from the back piece and attach onto the front. The collar needs to be secure in order to provide adequate support. It should be completely snug.

(See Pictures 3 and 4) Only once this step is complete can the other person stop holding the patient's head.

Picture 3




Picture 4



Once fitted correctly the chin should be centred and supported in the chin piece. No plastic should be touching the skin. The chin should not be able to be pushed down inside the collar, nor should this be attempted. (See Pictures 5 and 6)

Picture 5



Picture 6



Cleaning and skin care

Miami J Collar Basic Collar Care and Fitting Video

▶ <https://youtu.be/nanigBLYIP4>



- Keep the collar clean to prevent skin irritation.
- The pads should be changed if dirty or soiled. If you don't have a set of spare pads, contact Orthotics.
- It is easier to change one section of pads at a time to prevent confusion regarding the alignment of the pads. Before removing the soiled blue and grey pads look carefully at the shape and orientation of them so that you can reposition the clean pads properly.
- You can remove the blue pads from the white plastic shell and clean them with mild soap and water. Do not use bleach or harsh detergents. Thoroughly rinse the pads with clean water. Wring out the excess water and squeeze in a towel. Lay the pads flat and allow to air dry. You can wipe the white shell with soap and water and wipe it dry or let it air dry.
- Attach replacement pads, these should have been provided. Note: the blue side should be in contact with your skin ("blue is for you") and the grey side should contact the white plastic of the collar.
- Adjust the pads to make sure no plastic touches the skin.

Dos and don'ts while wearing the Miami J Collar

Do:

- Follow your Healthcare Professional's instructions about when and how long you should wear the collar.
- Have the collar tightened to maintain comfortable snugness.
- Minimise neck movements within the collar.

Don't:

- Stop using the collar unless your Healthcare Professional approves this.
- Take part in active sports.
- Engage in any physical activity that causes excessive pain.

Frequently Asked Questions

How long does the collar need to be worn for?

Most people wear the collar for up to 12 weeks after the injury. Your Consultant will tell you when you can stop wearing the collar, but you will likely be advised to wean out of it slowly rather than discarding it immediately.

DO NOT stop wearing the collar unless you have been advised to do so by a Healthcare Professional.

How comfortable should it be?

The collar is designed to restrict movement in the neck. This may feel uncomfortable at first and perhaps claustrophobic, but this should pass. If the skin on the chin becomes red or sore, try lying down for short periods or reclining in a comfortable chair to relieve the pressure.

You must not take the collar off at any time unless you are lying flat in bed and 2 people are present to hold your head and remove the collar for you.

What if the collar is too uncomfortable and I need to take it off?

If you have been wearing the collar without any problems up until now, first try to find out what could be causing the change in comfort.

Some common reasons are: pads moving, plastic digging in, sitting or lying in a different position, collar is not fastened tight enough or objects/food falling inside the collar.

If you continue to feel uncomfortable and feel you have to take the collar off, seek medical advice, but do not stop wearing the collar without speaking to a Healthcare Professional first as this increases the risk of permanent injury.

How do I dress myself?

While wearing the collar, you can dress in the normal way on sitting up out of bed. It's advisable to remain seated where possible to reduce the risk of falling over.

How do I bath or shower?

You must bath or shower with the collar on. Afterwards, you will need help from 2 people to change the collar pads over following the same instructions detailed earlier in this leaflet. The damp pads can be removed and left to air dry, while the fresh dry pads are put in the collar. Do not use heat to dry the pads as this may damage them. If you don't have a set of spare pads, contact Orthotics.

Can I remove the collar to eat?

No, you must keep your collar on while eating. If you find eating difficult in the collar, small changes can help such as cutting food up into smaller pieces or eating softer food and chewing more thoroughly. The collar can be cleaned after eating if necessary when you are lying down, again with help from 2 people if it needs to be removed.

When can I lift objects and what weight is OK?

Simple domestic objects are OK but avoid heavy weights. Avoid anything that puts a strain on your neck or which causes you discomfort. You should avoid any twisting or bending of your neck.

When can I return to physical activities?

Regular exercise is beneficial to a rapid recovery, and we recommend regular walking while in the collar or using an exercise bike, or gentle standing or seated exercises. Any exercise which involves a risk of falling should also be avoided.

Avoid any activity that puts strain on the neck or encourages attempts at movement within the collar. Some discomfort can be expected after starting to exercise. You should tell your Healthcare Professional about any new and excessive pain.

How far can I walk?

We recommend frequent short walks but be careful not to cause yourself pain. Most patients experience some muscle discomfort when they start regular activities. This should settle within 24 hours. If it continues check with a Healthcare Professional.

Can I drive while wearing my collar?

You should not drive while wearing the collar as your neck movement will be limited which will limit your line of vision and rotating your neck while wearing your collar can cause permanent injury. Fitness to drive is your personal responsibility and you should comply with the law and insurance requirements.

When can I travel or fly?

There are no medical restrictions on travelling or flying in a collar although there may be other medical reasons why it is not advisable to fly. You should check with your Healthcare Professional if this is something you are thinking about doing.

When can I return to work?

This will depend on the nature of your injury and job. Take advice from your Healthcare Professional as to when it is safe for you to return to your specific job.

Any problems?

If there are any problems with the collar you can contact one of the numbers below for advice or review as required.

Orthotics

☎ 0800 592087

✉ appointmentsbookingcentre@ggc.scot.nhs.uk

Or Contact the number below for the service that provided your collar.

Service/Location	
Healthcare Professional Contact	
Telephone Number	
Personalised Advice	

